




Draft Genome Sequence of the Plant Growth–Promoting *Pseudomonas punonensis* Strain D1-6 Isolated from the Desert Plant *Erodium hirtum* in Jordan

Feras F. Lafi,^{a,b} Maha L. AlBladi,^a Nida M. Salem,^c Luma Al-Banna,^c Intikhab Alam,^b Vladimir B. Bajic,^b Heribert Hirt,^a  Maged M. Saad^a

King Abdullah University of Science and Technology (KAUST), Biological and Environmental Sciences and Engineering Division (BESE), Thuwal, Saudi Arabia^a; King Abdullah University of Science and Technology (KAUST), Computational Bioscience Research Center (CBRC), Thuwal, Saudi Arabia^b; University of Jordan, Department of Agriculture, Amman, Jordan^c

ABSTRACT *Pseudomonas punonensis* strain D1-6 was isolated from roots of the desert plant *Erodium hirtum*, near the Dead Sea in Jordan. The genome of strain D1-6 reveals several key plant growth–promoting and herbicide-resistance genes, indicating a possible specialized role for this endophyte.

Herbicides are chemical compounds that are used in agriculture to fight off weeds and invading species of other noncrop plants. Recent discoveries have indicated that herbicides might be generated from microbial sources such as plant endophytes to protect plants from invasive and parasitic plant species (1, 2). Some of these endophytes produce antiherbicide properties that can counteract the function of commercial herbicides. Many of the species belong to the *Pseudomonas* genus (3, 4). In this study, we isolated *P. punonensis* strain D1-6 from surface-sterilized roots of the desert plant *Erodium hirtum* near the Dead Sea (31° 40.077 N; 35° 34.538E) in Jordan. Based on 16S rRNA gene analysis, strain D1-6 was closely related to *P. punonensis* (NR_109583) with 99% identity (5) and related to *P. fulva* (NR_074659) isolated from rice paddies. Genomic DNA of strain D1-6 was extracted using Qiagen’s DNeasy blood and tissue kit following the manufacturer’s protocol. The DNA library was constructed as described previously and then sequenced using paired-end reads by Illumina MiSeq (6).

MegaBLAST (7) searches of the strain D1-6 concatenated genome against the NCBI reference genome database (<http://www.ncbi.nlm.nih.gov/genome>) revealed that the closest relative genome belonged to *P. fulva* 12-X (CP002727) isolated from rice paddies with a query coverage of 77% and sequence similarity of 91%. We used SPAdes assembler version 3.6 with a 1-kb contig cutoff size (8) for contig assembly and then used the Indigo pipeline (9) for gene annotation and predicted open reading frames (ORFs) using FragGeneScan (10). *De novo* assembly of MiSeq reads for strain D1-6 resulted in 32 contigs with a total length of 4,534,589 bp and a mean contig size of 141,706 bp. The N_{50} was 322,232 bp and the L_{50} was reached in six contigs; the GC content of this genome was 61.4%. The annotation of *P. punonensis* strain D1-6 resulted in 3,244 ORFs, six rRNAs, 58 tRNAs, and 103 ncRNAs.

P. punonensis strain D1-6 produces the well-known plant growth–promoting enzyme 1-aminocyclopropane-1-carboxylate deaminase (ACC deaminase) (EC: 3.5.99.7, K01505) (11). Other plant growth–promoting enzymes such as cholesterol oxidase (EC: 1.1.3.6) provide insect resistance (12), and polygalacturonase (EC: 3.2.1.15) increases plant resistance to both fungal and bacterial pathogens via oligogalacturonide release (13). The presence of arginine decarboxylase (EC: 4.1.1.19) was shown to increase

Received 2 November 2016 Accepted 8 November 2016 Published 12 January 2017

Citation Lafi FF, AlBladi ML, Salem NM, Al-Banna L, Alam I, Bajic VB, Hirt H, Saad MM. 2017. Draft genome sequence of the plant growth–promoting *Pseudomonas punonensis* strain D1-6 isolated from the desert plant *Erodium hirtum* in Jordan. *Genome Announc* 5:e01437-16. <https://doi.org/10.1128/genomeA.01437-16>.

Copyright © 2017 Lafi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Heribert Hirt, heribert.hirt@kaust.edu.sa.

plant resistance to salinity, heat, and dehydration (14). *P. punonensis* strain D1-6 encodes for phosphinothricin acetyltransferase genes (EC: 2.3.1.183) that are responsible for herbicide resistance (15). The genome of *P. punonensis* strain D1-6 also encodes for many enzymes that have been identified as targets for the development of herbicide-resistant plants, e.g., quinate/shikimate dehydrogenase (EC: 1.1.1.282) (16), 4-hydroxyphenylpyruvate dioxygenase (EC: 1.13.11.27) (17). The finding of phosphinothricin (PPT) and other herbicide-resistance genes in *P. punonensis* strain D1-6 may indicate a possible role for this endophyte for providing the plant host with herbicide resistance.

Accession number(s). The genome of *Pseudomonas punonensis* D1-6 was deposited at DDBJ/EMBL/GenBank under the accession number [LWHA00000000](https://www.ncbi.nlm.nih.gov/nuclseq/LWHA00000000). The version described in this paper is the first version, LWHA01000000.

ACKNOWLEDGMENTS

Genome sequencing was performed at the biological core laboratories of the King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia. We are grateful for the use of the Dragon and Snap Dragon computer clusters at the Computational Bioscience Research Center (CBRC) at KAUST.

This work was supported by a base fund research grant to H.H. from King Abdullah University of Science and Technology (KAUST). The computational aspect of this work was supported by KAUST Office of Sponsored Research (OSR) under awards URF/1/1976-02 and FCS/1/2448-01 to V.B.B.

REFERENCES

- Harding DP, Raizada MN. 2015. Controlling weeds with fungi, bacteria and viruses: a review. *Front Plant Sci* 6:659. <https://doi.org/10.3389/fpls.2015.00659>.
- Li Y, Sun Z, Zhuang X, Xu L, Chen S, Li M. 2003. Research progress on microbial herbicides. *Crop Protect* 22:247–252. [https://doi.org/10.1016/S0261-2194\(02\)00189-8](https://doi.org/10.1016/S0261-2194(02)00189-8).
- Ramos JL, Duque E, Ramos-Gonzalez MI. 1991. Survival in soils of an herbicide-resistant *Pseudomonas putida* strain bearing a recombinant Tol plasmid. *Appl Environ Microbiol* 57:260–266.
- Moore JK, Braymer HD, Larson AD. 1983. Isolation of a *Pseudomonas* sp. which utilizes the phosphonate herbicide glyphosate. *Appl Environ Microbiol* 46:316–320.
- Ramos E, Ramírez-Bahena MH, Valverde A, Velázquez E, Zúñiga D, Velez-moro C, Peix A. 2013. *Pseudomonas punonensis* sp. nov., isolated from straw. *Int J Syst Evol Microbiol* 63:1834–1839. <https://doi.org/10.1099/ijs.0.042119-0>.
- Lafi FF, Bokhari A, Alam I, Bajic VB, Hirt H, Saad MM. 2016. Draft genome sequence of the plant growth-promoting *Cupriavidus gilardii* strain JZ4 isolated from the desert plant *Tribulus terrestris*. *Genome Announc* 4(4): e00678-16. <https://doi.org/10.1128/genomeA.00678-16>.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421. <https://doi.org/10.1186/1471-2105-10-421>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Alam I, Antunes A, Kamau AA, Ba Alawi WB, Kalkatawi M, Stingl U, Bajic VB. 2013. Indigo—integrated data warehouse of microbial genomes with examples from the Red Sea extremophiles. *PLoS One* 8:e82210. <https://doi.org/10.1371/journal.pone.0082210>.
- Rho MN, Tang HX, Ye YZ. 2010. FragGeneScan: predicting genes in short and error-prone reads. *Nucleic Acids Res* 38:e191. <https://doi.org/10.1093/nar/gkq747>.
- Loper JE, Hassan KA, Mavrodi DV, Davis EW, 2nd, Lim CK, Shaffer BT, Elbourne LD, Stockwell VO, Hartney SL, Breakwell K, Henkels MD, Tetu SG, Rangel LI, Kidarsa TA, Wilson NL, van de Mortel JE, Song C, Blumhagen R, Radune D, Hostetler JB, Brinkac LM, Durkin AS, Kluepfel DA, Wechter WP, Anderson AJ, Kim YC, Pierson LS III, Pierson EA, Lindow SE, Kobayashi DY, Raaijmakers JM, Weller DM, Thomashow LS, Allen AE, Paulsen IT. 2012. Comparative genomics of plant-associated *Pseudomonas* spp.: insights into diversity and inheritance of traits involved in multitrophic interactions. *PLoS Genet* 8:e1002784. <https://doi.org/10.1371/journal.pgen.1002784>.
- Pollegioni L, Piubelli L, Molla G. 2009. Cholesterol oxidase: biotechnological applications. *FEBS J* 276:6857–6870. <https://doi.org/10.1111/j.1742-4658.2009.07379.x>.
- Ferrari S, Galletti R, Pontiggia D, Manfredini C, Lionetti V, Bellincampi D, Cervone F, De Lorenzo G. 2008. Transgenic expression of a fungal *endo*-polygalacturonase increases plant resistance to pathogens and reduces auxin sensitivity. *Plant Physiol* 146:669–681. <https://doi.org/10.1104/pp.107.109686>.
- Liu JH, Ban Y, Wen XP, Nakajima I, Moriguchi T. 2009. Molecular cloning and expression analysis of an arginine decarboxylase gene from peach (*Prunus persica*). *Gene* 429:10–17. <https://doi.org/10.1016/j.gene.2008.10.003>.
- Páez-Espino AD, Chavarría M, de Lorenzo V. 2015. The two paralogue *phoN* (phosphinothricin acetyl transferase) genes of *Pseudomonas putida* encode functionally different proteins. *Environ Microbiol* 17:3330–3340. <https://doi.org/10.1111/1462-2920.12798>.
- Lindner HA, Nadeau G, Matte A, Michel G, Ménard R, Cygler M. 2005. Site-directed mutagenesis of the active site region in the quinate/shikimate 5-dehydrogenase YdiB of *Escherichia coli*. *J Biol Chem* 280: 7162–7169. <https://doi.org/10.1074/jbc.M412028200>.
- Falk J, Andersen G, Kernebeck B, Krupinska K. 2003. Constitutive over-expression of barley 4-hydroxyphenylpyruvate dioxygenase in tobacco results in elevation of the vitamin E content in seeds but not in leaves. *FEBS Lett* 540:35–40. [https://doi.org/10.1016/S0014-5793\(03\)00166-2](https://doi.org/10.1016/S0014-5793(03)00166-2).